

INSTITUUT VOOR PLANTENZIEKTENKUNDIG ONDERZOEK
WAGENINGEN, NEDERLAND
DIRECTEUR: Dr. J. G. TEN HOUTEN

MEDEDELING No. 267

**THE TRANSMISSION OF RATTLE VIRUS
BY TRICHODORUS PACHYDERMUS**

DOOR

H. H. SOL EN J. W. SEINHORST



OVERDRUK UIT:
T. PL.-ZIEKTEN, 67: 307-311, 1961

18 DEC 1961

INSTITUUT VOOR PLANTENZIEKTENKUNDIG ONDERZOEK (I.P.O.)

Office and main laboratory:

Binnenhaven 12, tel. 2151, 2152 en 3641
Wageningen, The Netherlands.

Staff:

Director:

Dr. J. G. TEN HOUTEN

Deputy director and head of the Entomological Dept.:

Dr. H. J. DE FLUITER, Wageningen.

Head of the Mycological Dept.:

Ir. J. H. VAN EMDEN, Wageningen.

Head of the Nematological Dept.:

Dr. Ir. J. W. SEINHORST, Wageningen.

Head of the Plant Disease Resistance Dept.:

Dr. J. C. s'JACOB, Wageningen.

Head of the Virological Dept.:

Miss Drs. F. QUAK, Wageningen.

Head of the Section Agricultural Aviation:

Miss M. C. KERSSSEN, Wageningen.

Head of the Section Biochemical Research and Application of Radioactive Isotopes:

Dr. J. H. VENKAMP, Wageningen.

Head of the Section Air Pollution Problems:

Ir. F. H. F. G. SPIERINGS, Wageningen.

Research workers at the Wageningen Laboratory:

Dr. Ir. A. B. R. BEEMSTER, Virologist

D. Z. MAAT, Virologist

Ir. J. H. VAN EMDEN, Phytopathologist

F. A. VAN DER MEER, Entomologist

Ir. J. A. DE BOKX, Virologist

Dr. J. C. MOOI, Phytopathologist

Dr. Ir. L. Bos, Virologist

J. P. W. NOORDINK, Radiochemist

Dr. H. H. EVENHUIS, Entomologist

W. C. NIJVELDT, Entomologist

Dr. H. J. DE FLUITER, Entomologist

Ir. H. DEN OUDEN, Nematologist

Dr. C. J. H. FRANSSEN, Entomologist

Miss Drs. H. J. PRAELTZER, Virologist

Dr. J. GROSJEAN, Phytopathologist

Ir. A. VAN RAAJ, Plantphysiologist

Ir. N. HUBBELING, Phytopathologist and plantbreeder

Miss Drs. F. QUAK, Virologist

Dr. J. C. s'JACOB, Phytopathologist and plantbreeder

Drs. L. E. VAN 'T SANT, Entomologist

Miss M. C. KERSSSEN, Agricultural aviation expert

Dr. Ir. J. W. SEINHORST, Nematologist

Miss Dr. C. H. KLINKENBERG, Nematologist

Dr. H. H. SOL, Virologist

Ir. R. E. LABRUYÈRE, Phytopathologist

Ir. J. VAN DER SPEK, Phytopathologist

Drs. H. P. MAAS GEESTERANUS, Phytopathologist

Ir. F. H. F. G. SPIERINGS, Plantphysiologist

G. M. TICHELAAR, Phytopathologist

Dr. F. TJALLINGII, Phytopathologist

Ir. E. UBELS, Phytopathologist

Dr. J. H. VENKAMP, Biochemist

Research workers elsewhere

Drs. J. M. M. v. BAKEL, Phytopathologist } detached to „Proefstation voor de Groenteteelt
Ir. C. KAAL, Nematologist } in de volle grond", Alkmaar, tel. 02200-6541.

Drs. D. J. DE JONG, Entomologist } detached to „Proefstation voor de Fruitteelt in de volle
Ir. G. S. ROOSJE, Phytopathologist } grond", Wilhelminadorp, Goes, tel. 01100-2261.

M. VAN DE VRIE, Entomologist }
Ir. T. W. LEFERING, Phytopathologist/Virologist, detached to „Proeftuin Noord Limburg"
Venlo, tel. 04700-2503.

Ir. F. A. HAKKAART, Virologist } detached to „Proefstation voor de bloemisterij
Drs. G. SCHOLTEN, Phytopathologist } in Nederland", Aalsmeer, tel. 02977-688.

Dr. K. VERHOEFF, Phytopathologist, detached to „Proeftuin voor de Groente en- Fruitteelt
onder glas", Naaldwijk, tel. 01740-4545.

Guest workers:

Dr. P. A. VAN DER LAAN, Entomologist, „Laboratorium voor toegepaste Entomologie der
Gemeente Universiteit", Amsterdam, tel. 020-56282.

Dr. Ir. G. S. VAN MARLE, Entomologist, Diepenveenseweg 226, Deventer, tel. 06700-3617.

Ir. G. W. ANKERSMIT, Entomologist, „Laboratorium voor Entomologie", Agricultural University, Wageningen, tel. 08370-2438.

Dr. Ir. J. B. M. VAN DINSTER, Entomologist, „Laboratorium voor Entomologie", Agricultural University, Wageningen, tel. 08370-2438.

Aphidological Adviser:

Mr. D. HILLE RIS LAMBERS, Entomologist, T.N.O., Bennekom, tel. 08379-2458.

THE TRANSMISSION OF RATTLE VIRUS BY *TRICHODORUS PACHYDERMUS*¹

Met een samenvatting:

De overdracht van ratelvirus door *Trichodorus pachydermus*

BY

H. H. SOL and J. W. SEINHORST

Instituut voor Plantenziektenkundig Onderzoek, Wageningen

HEWITT, RASKI and GOHEEN (1958) were the first to demonstrate the transmission of a soil-borne virus by nematodes. They were able to infect healthy grape vines with the fanleaf virus by placing in the soil near their roots the nematode *Xiphinema index* Thorne & Allen, taken from soil around the roots of diseased grape vines. *Xiphinema diversicaudatum* (Micoletzky) was later found to transmit *Arabis* mosaic virus (JHA & POSNETTE, 1959, 1961; HARRISON & CADMAN, 1959). The viruses transmitted by these *Xiphinema* species are isometric in shape.

Rod-shaped soil-borne viruses may also be transmitted by nematodes. SOL, VAN HEUVEN & SEINHORST (1960) were able to transmit rattle virus and *Atropa belladonna* mosaic virus to tobacco by growing it in soil to which had been added nematodes from soils infected with these viruses. *Hoplolaimus uniformis* Thorne and *Hemicycliophora* spec., isolated from infective bulk suspensions of nematodes, did not transmit the virus. Further experiments revealed that of four dorylaimoid species investigated, one, *Trichodorus pachydermus* Seinhorst, readily transmitted rattle virus.

MATERIALS AND METHODS

The nematodes used in the transmission experiments were extracted from sandy soil collected at Lisse (four dorylaimoid species including *Trichodorus pachydermus* of experiments A to D, listed in table 1) and at Gieterveen (experiment E in table 1). The Lisse field was known to be infected with rattle virus from observations by VAN SLOGTEREN (1958), which were corroborated in 1960 by the spinach test. For this test spinach is grown in the soil to be investigated. About five weeks after sowing, sap from the roots is pressed out and rubbed on leaves of healthy White Burley tobacco plants. The presence of rattle virus in the spinach roots, and therefore in the soil to be investigated, is revealed by the appearance of the necrotic ring and line patterns characteristic of rattle. In the field at Gieterveen potatoes suffered from stem-mottle and the presence of rattle virus could again be demonstrated by the spinach test.

Nematodes were extracted from soil by elutriation and sieving (SEINHORST, 1956). Not more than three to four hours after the elutriation specimens of the eelworm species to be investigated were picked from the extracted nema-

¹ Accepted for publication 15 June, 1961.

TABLE 1. The transmission of rattle virus by *Trichodorus pachydermus*.
De overdracht van ratelvirus door *Trichodorus pachydermus*.

Experiment <i>Proef</i>	Number of nematodes per bag <i>Aantal aaltjes per zakje</i>	Amount of soil per bag in g <i>Hoeveelheid grond per zakje in g</i>	Number of plants <i>Aantal planten</i>	Number of infected plants <i>Aantal geïnfecteerde planten</i>
A	50	125	5	2
	0	125	4	0
B	100	250	5	5
	0	250	5	0
C	10	50	7	4
	0	50	2	0
D	16	50	5	5
	8	50	10	10
	4	50	10	7
	2	50	10	7
	1	50	10	4
	0	50	5	0
E	8	50	7	7
	0	50	5	0
Total <i>Totaal</i>	inoculated soil <i>geïnoculeerde grond</i>		69	51
	controls <i>controles</i>		21	0

tode mixture and, before being transferred to the soil to be inoculated, placed in drops of distilled water on a watch glass. The soil used in the experiments was autoclaved and, in order to remove water-soluble substances which might inactivate the nematodes, hot water was sucked through the soil on a Buchner funnel. From 50 to 250 g of this soil was placed in a plastic bag of suitable size. The nematodes were transferred to it with a little water and not more than a day afterwards one seedling of *Nicotiana tabacum* var. White Burley was planted in each bag. In the tests to investigate which nematode species transmits rattle, at least thirty specimens of one species were placed in a bag.

Transmission of virus was sometimes revealed by symptoms in the leaves of the test plants but more often had to be determined by testing for the presence of virus in their roots. Presence or absence of virus in the roots of control plants and of plants which showed no leaf symptoms after having grown for two to three weeks in soil inoculated with nematodes was investigated as described for the spinach test.

RESULTS

Of the four dorylaimoid species tested, rattle virus was transmitted only by *Trichodorus pachydermus*. Experiment D of table 1 shows that even one specimen of this species can infect a tobacco plant. That not all specimens added singly to the bags transmitted virus may have been due to the fact that not all

of them carried virus, or possibly to inactivity or death of some of the specimens. Generally no more than about two out of every three specimens added to soil could be extracted from it a week later, indicating that approximately one in three of the nematodes became inactive after transfer.

GEOGRAPHICAL DISTRIBUTION OF RATTLE VIRUS AND OF *TRICHODORUS PACHYDERMUS*

T. pachydermus occurs quite commonly on sandy soils in the Netherlands whether arable land, pasture, woodland, or soil which is not under cultivation. It seems to be rare in clay soils, as is the soil-borne rattle virus. The distribution of the virus may therefore very well be limited to areas, where *T. pachydermus* occurs. This must still be corroborated by further research.

SAMENVATTING

De nematode-soort *Trichodorus pachydermus* Seinhorst kan ratelvirus overbrengen naar gezonde White Burley tabaksplanten. Eén aaltje, geïsoleerd uit met ratelvirus besmette grond, is zelfs in staat een tabaksplant met het virus te infecteren.

Evenals ratelvirus komt deze nematode-soort vooral voor in lichtere grondsoorten. In kleigrond worden beide zelden aangetroffen. Het is mogelijk, dat de verspreiding van het virus beperkt is tot gebieden waar *T. pachydermus* voorkomt. Dit moet nog nader worden onderzocht.

ACKNOWLEDGEMENTS

The authors are greatly indebted to Ir. D. H. M. VAN SLOGTEREN, Laboratorium voor Bloembollenonderzoek, Lisse and Dr. Ir. E. G. KLOOSTERMAN, N.A.K., Wildervank for their help in locating infected fields and collecting soil samples.

REFERENCES

- HARRISON, B. D. & C. H. CADMAN, - 1959. Role of a dagger nematode (*Xiphinema* sp.) in outbreaks of plant diseases caused by *Arabis* mosaic virus. *Nature* 184: 1624-1626.
- HEWITT, W. B., D. J. RASKI & A. C. GOHEEN, - 1958. Nematode vector of soil-borne fanleaf virus. *Phytopathology* 48: 586-595.
- JHA, A. & A. F. POSNETTE, - 1959. Transmission of a virus to strawberry plants by a nematode (*Xiphinema* sp.). *Nature* 184: 962-963.
- JHA, A. & A. F. POSNETTE, - 1961. Transmission of *Arabis* mosaic virus by the nematode *Xiphinema diversicaudatum* (Micol.). *Virology* 13: 1-12.
- SEINHORST, J. W., - 1956. The quantitative extraction of nematodes from soil. *Nematologica* 1: 249-268.
- SLOGTEREN, D. H. M. VAN, - 1958. Ratievirus als oorzaak van ziekten in bloembolgewassen en de mogelijkheden de infectie door middel van grondontsmetting te bestrijden. *T.Pl.-ziekten* 64: 452-462.
- SOL, H. H., J. C. VAN HEUVEN & J. W. SEINHORST, - 1960. Transmission of rattle virus and *Atropa belladonna* mosaic virus by nematodes. *T.Pl.-ziekten* 66: 228-231.

**Mededelingen van het Instituut voor Plantenziektenkundig Onderzoek
verschenen in de tweede helft van 1961**

**Contributions of the Institute for Phytopathological Research
published during the second half of 1961**

Binnenhaven 12, Wageningen, The Netherlands

- No 255. FRANSSEN, C. J. H. en W. P. MANTEL, De door tripsen veroorzaakte beschadigingen in het vlasgewas en het voorkómen daarvan. (Preventing damage by thrips in flax.) T.Pl.-ziekten, 67: 39-51, 1961. Prijs no. 255 f 0,50.
- No 256. ZADOKS, J. C., Yellow rust on wheat. Studies in epidemiology and physiologic specialisation. T.Pl.-ziekten, 67: 69-256, 1961. Prijs no. 256 f 3,60.
- No 257. BOKX, J. A. DE, Waardplanten van het aardappel-Y^N-virus. (Host plants of the potato virus Y^N (tobacco vein necrosis virus)) T.Pl.-ziekten, 67: 273-277, 1961. Prijs no. 257 f 0,25.
- No 258. BEEMSTER, A. B. R., Translocation of leaf roll and virus Y in the potato. Proc. 4th Conf. Potato Virus Dis. Braunschweig, 1960: 60-67, 1961. Prijs no. 258 en 259 samen f 0,60.
- No 259. BEEMSTER, A. B. R., Een vergelijking tussen het transport van X-virus en twee verschillende stammen van Y-virus in aardappelplanten. (A comparison between the translocation of virus X and two different strains of virus Y in the potato plant). T.Pl.-ziekten, 67: 278-279, 1961. Prijs no. 258 en 259 samen f 0,60.
- No 260. FLUITER, H. J. DE, Die nützlingsschonende Schädlingbekämpfung und ihre Beziehung zur Taxonomie der Nützlinge und Schädlinge. Entomophaga, VI: 143-154, 1961. Prijs no. 260 f 0,35.
- No 261. EVENHUIS, H. H. en D. J. DE JONG, Oecologische waarnemingen over bladluizen van appel in 1960. (Ecological observations on apple aphids in 1960). Meded. Dir. Tuinbouw, 24: 285-290, 1961. Prijs no. 261 f 0,30.
- No 262. ROOSJE, G. S. en C. A. R. MEIJNEKE, 'Fire blight', een dreigend gevaar voor de Nederlandse fruitteelt? (Fire blight: an imminent danger to fruit-growing in The Netherlands). Meded. Dir. Tuinbouw, 24: 356-365, 1961. Prijs no. 262 f 0,55.
- No 263. BOS, L. en J. P. H. VAN DER WANT, Legume virus research in The Netherlands with special reference to forage crops. Proc. 8th int. Grassland Congr., 1960: 207-210, 1961. Prijs no. 263 f 0,25.
- No 264. TICHELAAR, G. M., De invloed van Gladiolus op de kieming van sclerotiën van Sclerotium cepivorum. (The influence of Gladiolus on the germination of sclerotia of Sclerotium cepivorum). T.Pl.-ziekten, 67: 290-295, 1961. Prijs no. 264 f 0,45.
- No 265. BAKKER, M. en G. SCHOLTEN, Bacterieziekten bij Amerikaanse anjers in Nederland. (Bacterial diseases of carnations in The Netherlands.) T.Pl.-ziekten, 67: 296-302, 1961. Prijs no. 265 f 0,30.
- No 266. BOKX, J. A. DE, Een vergelijking tussen verschillende toetsmethoden voor het aardappel-Y-virus. (A comparison of different methods for testing potato virus Y.) T.Pl.-ziekten, 67: 303-306, 1961. Prijs no. 266 f 0,25.
- No 267. SOL, H. H. en J. W. SEINHORST, The transmission of rattle virus by Trichodorus pachydermus. T.Pl.-ziekten, 67: 307-311, 1961. Prijs no. 267 f 0,25.

